

Global sensitivity analysis for models described by continuous time Markov chains with application to epidemic models

H. M. KOUYE
Paris-Saclay University

Supervisor(s): Gildas Mazo (MaIAGE, INRAE), Elisabeta Vergu (MaIAGE, INRAE) and Clémentine Prieur (LJK, Grenoble-Alpes University)

PhD expected duration: Oct. 2019 - Sep. 2022

Address: UR1404 INRAE MaIAGE (Mathématiques et Informatique Appliquées du Génome à l'Environnement), 78352 Jouy-en-Josas

Email: henri-mermoz.kouye@inrae.fr

Abstract: Stochastic models are increasingly used in various fields (epidemiology, biology, physics etc) to describe different phenomena. They can provide insights into complex situations at large to small scales. Models can be described by stochastic processes, hence including an intrinsic randomness. Performing global sensitivity analysis (GSA) for such models is challenging, if the intrinsic randomness of the system is considered as a noise on specific quantities of interests (QoIs). The objective of our work is to propose a generic strategy to perform GSA on such stochastic systems. In this context, several studies have been led to point out the impact of the inputs on the stochastic QoIs. Global sensitivity analysis methods (Sobol', Polynomial chaos expansion, Karhunen-Loeve decomposition, ...) have been implemented for stochastic models with scalar QoIs [4], [7], models described by stochastic differential equations [2], Markov chain based models for chemical physics [8], etc. However, either these studies did not propose a generic framework Markov chain based models or they only deal with scalar QoIs.

In order to introduce a more generic framework, our strategy aims firstly at separating the two sources of variability, namely parameters uncertainty and intrinsic randomness and secondly at putting the model functional QoI Y under the form:

$$Y = f(X, Z), \tag{1}$$

where f is a function, X stands for the uncertain parameters and Z represents the intrinsic randomness such as X and Z are independent. For this purpose, two approaches are introduced: Sellke construction [9] and Kurtz representation [1],[5]. Depending on the class of models, Sellke construction allows to describe Z as a finite-dimensional vector with known distribution while under Kurtz representation, Z is a vector of independent Poisson processes with intensity 1. These two approaches allow not only to estimate the sensitivity indices of the input parameters but also to measure the influence of the intrinsic randomness on the global variability of the QoI under study. Furthermore, Sellke construction can be used to extend to the non-markovian framework.

From (1), we put ourselves in various frameworks of sensitivity analysis and implement different methods (global sensitivity indices [6], general metric space sensitivity indices [3], etc) by introducing the indices and their estimators and finally by carrying out the simulations. In order to illustrate the approaches we developed, they will be applied to some SIR-like models (fig. 1) which are usually used in stochastic modeling of epidemics.




Figure 1: Example of the SEIR stochastic epidemic model

References

- [1] Stewart N. Ethier and Thomas G. Kurtz. *Markov processes – characterization and convergence*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1986.
- [2] Pierre Etoré, Clémentine Prieur, Dang Khoi Pham, and Long Li. Global sensitivity analysis for models described by stochastic differential equations. *Methodology and Computing in Applied Probability*, pages 1–29, 2019.
- [3] Fabrice Gamboa, Thierry Klein, Agnès Lagnoux, and Leonardo Moreno. Sensitivity analysis in general metric spaces. working paper or preprint, July 2020.
- [4] J. L. Hart, A. Alexanderian, and P. A. Gremaud. Efficient computation of sobol' indices for stochastic models. *SIAM Journal on Scientific Computing*, 39(4):A1514–A1530, 2017.
- [5] Thomas G. Kurtz. Representation and approximation of counting processes. In Wendell H. Fleming and Luis G. Gorostiza, editors, *Advances in Filtering and Optimal Stochastic Control*, pages 177–191, Berlin, Heidelberg, 1982. Springer Berlin Heidelberg.
- [6] Matieyendou M. Lamboni, Herve H. Monod, and David D. Makowski. Multivariate sensitivity analysis to measure global contribution of input factors in dynamic models. *Reliability Engineering and System Safety*, 96(4):450–459, 2011.
- [7] Gildas Mazo. An optimal tradeoff between explorations and repetitions in global sensitivity analysis for stochastic computer models. July 2019.
- [8] M. Navarro Jimenez, O. P. Le Maître, and O. M. Knio. Global sensitivity analysis in stochastic simulators of uncertain reaction networks. *The Journal of Chemical Physics*, 145(24):244106, December 2016.
- [9] Thomas Sellke. On the asymptotic distribution of the size of a stochastic epidemic. *Journal of Applied Probability*, 20(2):390–394, 1983.

Short biography – After a background in applied and fundamental mathematics, I obtained a master’s degree in mathematics applied to life sciences at the Paris-Saclay University. I am currently pursuing a thesis thanks to an INRAE-Inria funding and my work aims at studying sensitivity analysis methods for stochastic models with functional outputs.